

UNCLASSIFIED

AD NUMBER
AD464209
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; May 1965. Other requests shall be referred to Commanding Officer, US Army Biological Laboratories, Attn: TRB/TID, Fort Detrick, MD 21701.
AUTHORITY
BDRL, D/A ltr, 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

TECHNICAL MANUSCRIPT 223

COMPLEX AND CHEMICALLY DEFINED MEDIA
FOR THE GROWTH
OF CRYPTOCOCCUS NEOFORMANS

Ernest M. Snyder
John L. Converse
Norman D. Gary

MAY 1965

NOT TO BE REPRODUCED
WITHOUT PERMISSION
EXCEPT BY USERS OF THE

JUN 11 1965

UNITED STATES ARMY
BIOLOGICAL LABORATORIES
FORT DETRICK

BEST AVAILABLE COPY

20041122050

CATALOGED BY: DDC

AS AM M7

464209

AD464209

Reproduction of this publication in whole or part is prohibited except with permission of Commanding Officer, U.S. Army Biological Laboratories, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

Qualified requestors may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

900411020

605494

U.S. ARMY BIOLOGICAL LABORATORIES
Fort Detrick, Frederick, Maryland

TECHNICAL MANUSCRIPT 223

COMPLEX AND CHEMICALLY DEFINED MEDIA FOR
THE GROWTH OF CRYPTOCOCCUS NEOFORMANS

Ernest M. Snyder

John L. Converse

Norman D. Gary

Medical Bacteriology Division
DIRECTORATE OF BIOLOGICAL RESEARCH

Project 16522301A05901

May 1965

ABSTRACT

A maltose, ammonium acetate, inorganic salts medium containing thiamine and glutamate was formulated for growth of Cryptococcus neoformans; from which growth levels of approximately 10^8 cells per ml could be obtained within 72 to 96 hours incubation. A complex medium (nutrient broth with maltose and thiamine) with comparable growth characteristics, was also recommended. Both media supported very well the growth of 11 of 13 strains tested.

COMPLEX AND CHEMICALLY DEFINED MEDIA FOR THE GROWTH OF CRYPTOCOCCUS NEOFORMANS

Previous studies on the growth of Cryptococcus neoformans characterized the carbon and nitrogen assimilation spectra of the organism, its temperature tolerance, its aerobic nature, and its requirement for thiamine.¹⁻⁴ The majority of those studies employed solid media. In work employing liquid media, viable cell yields, when reported, have been low ($<1 \times 10^7$ /ml) even with extended incubation periods of 7 to 8 days. This manuscript describes chemically defined and complex media yielding viable counts in excess of 1×10^8 /ml after 48 to 72 hours' incubation. The recommended compositions of the media are given in Table 1.

TABLE 1. RECOMMENDED MEDIA FOR GROWTH
OF CRYPTOCOCCUS NEOFORMANS^{a/}

Constituent	Concentration, grams per liter
<u>Complex Medium</u>	
Maltose	40
Bacto-Peptide	5
Bacto-Beef Extract	3
Thiamine-HCl	100 µg/ml
<u>Chemically Defined Medium</u>	
Maltose	40
Ammonium Acetate	2
Sodium Glutamate	2
MgSO ₄ ·7H ₂ O	1.97
ZnSO ₄ ·7H ₂ O	0.0176
KH ₂ PO ₄	2.04
K ₂ HPO ₄	2.61
Thiamine-HCl	1 µg/ml

a. Initial pH: approximately 6.8 for each medium.

Results of growth studies employing complex medium are summarized in Table 2. Initial experiments using strain C-1, with a basal medium (0.5% peptone, 0.3% beef extract, 2% glucose, and thiamine-HCl), confirmed the requirements for aeration (shaken cultures) and thiamine (100 µg/ml). Inoculum size, over a range of 3×10^4 to 220×10^4 cells per 50 ml of medium, was not critical for maximum growth. Moreover, significantly higher cell yields were obtained at 25 C than at 34 C. Examination of various carbon sources established maltose (4%) as the best substrate, yielding approximately 1×10^9 cells per ml. The maximum growth was limited (30% less) by buffering the medium (pH 7.0); the culture pH at optimal growth was approximately 4.0.

Results of growth studies employing chemically defined medium are also shown in Table 2. Initial growth studies (strain C-1) compared various combinations of the constituents of the ammonium sulfate medium of Littman⁶ and the ammonium acetate medium of Roessler et al.⁶ The addition of thiamine and/or sodium glutamate (at the concentrations specified in the ammonium sulfate medium) to the ammonium acetate medium resulted in equal or better growth (approximately 2×10^8 cells per ml). The substitution of maltose for glucose in the ammonium acetate medium increased the growth level to 6×10^8 cells per ml. Supplements of the calcium, iron, manganese, or molybdenum salts to Littman's medium failed to improve the ammonium acetate medium. The substitution of ammonium chloride or ammonium sulfate, either singly or in combination, resulted in only 30 to 50% of the growth obtained with ammonium acetate.

Using 4% maltose as the energy source and decreasing the ammonium acetate level from 0.6% to 0.2% resulted in growth in excess of 1×10^9 cells per ml (at 48 hours' incubation). Increasing the level of sodium glutamate to 1 to 2% resulted in a slight increase in growth, but even at the 4 to 6% level, glutamate did not serve as a sole source of energy in the absence of maltose. One µg/ml of thiamine was sufficient for optimal growth. The original levels of magnesium and zinc in the ammonium acetate medium (0.2% and 0.00176%, respectively) were optimal for highest growth. Deletion experiments with the complete medium showed no decrease in growth at one week by omission of sodium glutamate, zinc sulfate, or dipotassium phosphate, but 20 to 30% growth reduction occurred during earlier periods of incubation (48 to 72 hr).

Growth of 13 strains of *C. neoformans*, in the chemically defined and complex media, is shown in Table 2. The majority of these strains (8 of 13) demonstrated higher growth in the chemically defined than in the complex medium. Considering both media, growth of 12 of the 13 strains approached 10^9 cells per ml or better.

TABLE 2. VIABLE CELL YIELDS OF VARIOUS STRAINS OF C. NEOFORMANS
IN COMPLEX AND CHEMICALLY DEFINED MEDIA^{a/}

Strain	Complex Medium		Chemically Defined Medium	
	48 hr	72 hr	48 hr	72 hr
C-1 ^{b/}	87	103	121	128
Scott ^{c/}	69	75	59	84
Kresge ^{c/}	86	80	65	74
Gibson ^{c/}	93	83	120	108
Green ^{c/}	- ^{e/}	31	-	3
Tucker ^{c/}	74	72	72	86
Bishop ^{c/}	68	65	88	120
Stanford ^{c/}	140	158	207	232
Stokes ^{c/}	96	88	105	116
32 ^{d/}	-	101	-	6
34 ^{d/}	-	69	-	56
35 ^{d/}	-	128	-	89
39 ^{d/}	-	58	-	85

a. Mean values of duplicate cultures expressed as 10^7 viable cells per ml.

b. From Dr. Libero Ajello, CDC, Atlanta, Georgia.

c. From Dr. C.W. Emmons, NIH, Bethesda, Md.

d. From Dr. Milton Huppert, VA Hospital, San Fernando, Calif.

e. Not counted.

LITERATURE CITED

1. Kuhn, L.R. 1939. Growth and viability of Cryptococcus hominis at mouse and rabbit body temperature. Proc. Soc. Exp. Biol. Med. 41:573-574.
2. Reid, J.D. 1949. The influence of vitamin B complex on the growth of Torulopsis (Cryptococcus) neoformans on a synthetic medium. J. Bacteriol. 58:777-782.
3. Evans, E.E., and E.R. Harrell. 1952. Cryptococcosis (Torulosis): A review of recent cases. Univ. of Mich. Med. Bull. 18:43-63.
4. Littman, M.L. 1957. An improved method for detection of urea hydrolysis by fungi. J. Infect. Dis. 101:51-61.
5. Littman, M.L. 1958. Capsule synthesis by Cryptococcus neoformans. Trans. N.Y. Acad. Sci. 20(7):623-648.
6. Roessler, W.G., E.J. Herbst, W.G. McCullough, R.C. Mills, and C.R. Brewer. 1946. The nutrition of Coccidioides immitis in submerged culture. J. Bacteriol. 51:572.

DISTRIBUTION LIST

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Assistant Scientific Director Building 812	1
Directorate of Biological Research Building 560	1
Directorate of Development Building 824	1
Directorate of Industrial Health & Safety Building 550	1
Chief, Program Coordination Office Building 812	1
Chief, Aerobiology Division Building 459	1
Chief, Medical Bacteriology Division Building 560	10
Chief, Medical Investigations Division Building 604	1
Chief, Physical Sciences Division Building 568	1
Chief, Process Development Division Building 469	1
Chief, Technical Evaluation Division Building 568	1
Chief, Virus & Rickettsia Division Building 539	1
Chief, Biomathematics Division Building 1422	1
Documents, Technical Library Building 426	2
Test Chamber Branch Technical Evaluation Division Building 1412	1

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Technical Releases Branch Technical Information Division Building 426	10
Editorial Branch Technical Information Division Building 816	1
U.S. Army Medical Unit Ft. Detrick, Maryland Building 120	1
Liaison Representative/Animal Disease Investigations Building 1301	5
U.S. Public Health Service Liaison Office Building 1301	9
Commanding Officer U.S. Naval Unit Building 125	3
Commanding General U.S. Army Edgewood Arsenal ATTN: SMUEA-CS Edgewood Arsenal, Maryland, 21010	1
Commanding Officer U.S. Army Chemical Research & Development Laboratories ATTN: Librarian Edgewood Arsenal, Maryland, 21010	2
Commanding General U.S. Army Munitions Command ATTN: AMSMU-SS Dover, New Jersey, 07801	1
Commanding General U.S. Army Munitions Command ATTN: AMSMU-RE-R Dover, New Jersey, 07801	1
Commandant U.S. Army CBR Weapons Orientation Course Dugway Proving Ground Dugway, Utah, 84022	1

ADDRESSEENUMBER OF COPIES

Commanding General
Deseret Test Center
ATTN: Technical Library
Fort Douglas, Utah, 84113

2

Commanding General
U.S. Army Materiel Command
Research Division, AMCRD-RC
R&D Directorate
Washington, D.C., 20315

1

Department of the Army
Asst Chief of Staff/Force Operations
ATTN: Technical Coordinator (B)
CBR & N Directorate
The Pentagon
Washington 25, D.C.

1

Defense Documentation Center
Cameron Station
Alexandria, Virginia, 22314

20

Detachment 4, RTD (ATCB)
Eglin Air Force Base, Florida, 32542

1

APGC (PGBAP-1)
Eglin Air Force Base, Florida, 32542

1

Dr. S.H. Madin
Scientific Director
Naval Biological Laboratory
Naval Supply Center
Oakland, California, 94614

1

Commander (Code 4036)
U.S. Naval Ordnance Test Station
China Lake, California, 93557

1

Commanding Officer and Director
U.S. Naval Applied Science Laboratory
Naval Base, Code 9440
Brocklyn, New York, 11251

1

U.S. Army Medical R&D Command
Office of the Surgeon General
ATTN: MEDDH-C
Main Navy Building, Room 2526
Washington, D.C., 20315

1

ADDRESSEENUMBER OF COPIES

Commandant
USA CmlCen & Sch, ATTN: Bio Branch
Ft. McClellan, Alabama, 36205

2

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is to be effected)		
1. ORIGINATING ACTIVITY (Corporate author) U.S. Army Biological Laboratories Fort Detrick, Frederick, Maryland, 21701		2a. REPORT SECURITY CLASSIFICATION <u>Unclassified</u> 2b. GROUP
3. REPORT TITLE COMPLEX AND CHEMICALLY DEFINED MEDIA FOR THE GROWTH OF <u>CRYPTOCOCCUS NEOFORMANS</u>		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Snyder, Ernest M. Gary, Norman D. Converse, John L.		
6. REPORT DATE May 1965	7a. TOTAL NO. OF PAGES 14 pages	7b. NO. OF REFS 6
8a. CONTRACT OR GRANT NO. b. PROJECT NO. c. IC522301A05901 d.	9a. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 223 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY LIMITATION NOTICES Qualified requestors may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY U.S. Army Biological Laboratories Fort Detrick, Frederick, Maryland, 21701	
13. ABSTRACT A maltose, ammonium acetate, inorganic salts medium containing thiamine and glutamate was formulated for growth of <u>Cryptococcus neoformans</u> ; from which growth levels of approximately 10^7 cells per ml could be obtained within 72 to 96 hours incubation. A complex medium (nutrient broth with maltose and thiamine) with comparable growth characteristics, was also recommended. Both media supported very well the growth of 11 of 13 strains tested.		

DD FORM 1473

Unclassified

Security Classification